DEPARTMENT of ENVIRONMENTAL SERVICES Water Supply & Pollution Control Division - Biology Bureau

LAKE TROPHIC DATA

MORPHOMETRIC:

Lake: SWANZEY LAKE	Lake Area (ha):	47.31
Town: SWANZEY	Maximum depth (m):	16.2
County: Cheshire	Mean depth (m):	6.9
River Basin: Connecticut	Volume (m³):	3271500
Latitude: 42°51' N	Relative depth:	2.0
Longitude: 72°18′ W	Shore configuration:	1.39
Elevation (ft): 524	Areal water load (m/yr)	3.94
Shore length (m): 3400	Flushing rate (yr ⁻¹):	0,60
Watershed area (ha): 414.4	P retention coeff.:	0.70
% watershed ponded: 0.0	Lake type:	natural

15 January 1987	4 August 1986
ASTERIONELLA 67%	CHRYSOSPHAERELLA 45%
TABELLARIA 22%	DINOBRYON 30%
	ASTERIONELLA 15%
	965.0
	3.93
KELLICOTTIA 80%	NAUPLIUS LARVAE 34%
NAUPLIUS LARVAE 10%	KERATELLA 17%
KERATELLA 4%	BOSMINA 14%
529	50
76	146
610	198
	Scattered
	4.5
3.6	0.2
	10
	< 10
	ASTERIONELLA 67% TABELLARIA 22% KELLICOTTIA 80% NAUPLIUS LARVAE 10% KERATELLA 4% 529 76 610

SUMMER THERMAL STRATIFICATION:

stratified

Depth of thermocline (m): 6.0 Hypolimnion volume (m³): 460500

CHEMICAL:	Lake: SWANZEY LAKE Town: SWANZEY				
	15 January 1987		4 August 1986		
DEPTH (m)	5.0	10.0	2.0	8.0	13.0
pH (units)	6.6	6.5	7.0	6.8	6.4
A.N.C. (Alkalinity)	5.7	6.0	5.3	5.4	7.9
NITRATE & NITRITE NITROGEN	0.33	< 0.05	< 0.05		< 0.05
TOTAL KJELDAHL NITROGEN	0.38	0.40	0.55	0.50	0.45
TOTAL PHOSPHORUS	0.008	0.015	0.004	0.011	0.019
CONDUCTIVITY (p mhos/cm)	38.7	37.6	36.0	35.7	38.8
APPARENT COLOR (cpu)	15	16	5	10	25
MAGNESIUM			0.65		
CALCIUM			2.8		
SODIUM			2.2		
POTASSIUM			0.60		
CHLORIDE	2	3	3		3
SULFATE					
TN : TP	89	27	138		24
CALCITE SATURATION INDEX			3.1		


All results in mg/L unless indicated otherwise

TROPHIC CLASSIFICATION: 1986

D.O.	S.D.	PLANT	CHL	TOTAL	CLASS
6	1	1	0	8	Meso.

COMMENTS:

- 1. The maximum depth on the sounding map is 52 feet (15.8m); however, a depth of 16.2 meters was found during the winter sampling.
- 2. This lake was previously surveyed in 1977. It received 3 more trophic points in 1986 than in 1977, and that was enough to move it from oligotrophic to mesotrophic. The major change during the 10 year period was an increase in weed growth.
- 3. The whole-water phytoplankton was 35% blue-greens, 30% cryptomonads, and 20% greens. Dominant genera were Chroococcus (30%) and Chroomonas (20%).



FIELD DATA SHEET

LAKE: SWANZEY LAKE TOWN: SWANZEY

DATE: 08/04/86 WEATHER: FAIR, PARTLY CLOUDY


DEPTH (M)	TEMP (°C)	*DISSOLVED OXYGEN	OXYGEN SATURATION
0.1	25.8	8.5	100 %
1.0	25.0	8.0	95 %
2.0	24.8	7.1	83 %
3.0	24.7	6.5	76 %
4.0	24.2	6.5	76 %
5.0	23.5	6.5	74 %
6.0	19.1	8.0	86 %
7.0	14.5	9.0	87 %
8.0	11.8	9.0	81 %
9.0	9.2	7.2	62 %
10.0	8.0	3.8	31 %
11.0	7.1	0.5	4 %
12.0	7.0	0.0	0 %
13.0	6.8	0.2	2 %
14.0	6.6	0.2	2 %
15.0	6.6	0.2	2 %

SECCHI DISK (m): 4.5 COMMENTS:

BOTTOM DEPTH (m): 15.5

TIME: 1240

*Dissolved oxygen values are in mg/L



AQUATIC PLANT SURVEY

LAKE: SWANZEY LAKE TOWN: SWANZEY DATE: 08/04/86

Key	PL.	ABUNDANCE	
ney	GENERIC	COMMON	ABOINDANCE
Ε	Eriocaulon septangulare	Pipewort	Scattered
Χ		Bottom growth	Scattered
U	Utricularia	Bladderwort	Sparse
Ь	Scirpus	Bulrush	Sparse
f	Chlorophyceae	Filamentous green algae	Scattered
С	Cyperaceae	non-flowering sedge	Sparse
N	Nymphaea	White water lily	Sparse
Υ	Nuphar	Yellow water lily	Sparse
W	Potamogeton	Pondweed	Sparse
Р	Pontederia cordata	Pickerelweed	Sparse

OVERALL ABUNDANCE: Scattered

GENERAL OBSERVATIONS:

- 1. Pipewort was relatively common around most of the shoreline, but because of its sparse growth habit, was listed as scattered.
- 2. Plants were locally abundant at the outlet area (locally called the Mill Pond), and at the northeast end near the inlet.
- 3. Filamentous algae and bottom growth were common along much of the shoreline.